



# BOTSWANA

## ABOUT BOTSWANA VACCINATIONS

Botswana, a landlocked southern African country, occupies the centre of the South African plateau.. Botswana's subtropical climate features an extended winter dry season from May through August. Toward the end of winter, monsoon winds bring rain, with eastern regions receiving extended 'rains' and the heaviest rainfall. June is the coolest month, October the warmest. In the eastern highlands, the July-Oct average temp ranges from 11c to 18c, in lower areas 14c to 21c.

### OVERALL RISK

High Risk - Preventable disease is much more common than in Australia. Consult a Travelvax clinic for your specific risk.

### MALARIA RISK

Malaria is present throughout country. Anti-malarial medications recommended. Type of malaria tablets will vary according to itinerary, length of stay and needs to be discussed with a medical practitioner.

## VACCINE PREVENTABLE DISEASES

### YELLOW FEVER

A yellow fever vaccination certificate is required for travellers aged 1 year or over arriving from or having transited through countries with risk of yellow fever transmission.

No vaccination certificate is required for direct travel from Australia or New Zealand.

### COVID-19

For the most up to date requirements for entry into Australia, go to [Smartraveller](#).

Please click this [COVID-19 Travel Regulations Map](#) (IATA) to view the latest Botswana COVID travel requirements.

### HEPATITIS A

Risk of Hepatitis A infection is high for all travellers, vaccination recommended.

### HEPATITIS B

Moderate presence of Hepatitis B in local population. Vaccination would be recommended.

## TYPHOID

Moderate risk for most travellers. Vaccination recommended for travel to smaller cities, villages and rural areas outside usual tourist routes. Some medical conditions pre-dispose to infection; whether vaccinations would be recommended should be discussed with a medical practitioner. Consult a medical practitioner for your specific risk.

## INFLUENZA

Year round risk present. Most common vaccine preventable illness in travellers. Vaccine recommended, effective for 1 year.

## RABIES

Disease present. Recommendation for vaccination will depend on specific itinerary and activities planned. Generally rabies vaccination is advised for high risk individuals such as veterinarians or animal handlers, cavers. Additionally for higher risk travellers who plan: extended periods outdoors, rural travel, adventurous activities including bicycling; also expats or long-term travellers to endemic regions and children (risk of more severe or risk-prone bites and may not report contact at all).

## JAPANESE ENCEPHALITIS

No risk to travellers.

## POLIO

No risk to travellers.

## TICKBORNE ENCEPHALITIS

No risk to travellers.

## MENINGOCOCCAL MENINGITIS

Low risk to travellers.

## CHOLERA

No risk to travellers.

## OTHER DISEASES

### TRAVELLERS DIARRHOEA

High risk. Travellers' Diarrhoea affects 20-60% of overseas travellers, food and water precautions are essential. Self-treatment medications may be recommended.

### INSECT-BORNE DISEASES

Diseases such as [Dengue](#), [Chikungunya](#), [Zika](#) may be present. Seasonal risk will vary by country. Whether preventive measures will be recommended will depend on itinerary, length of stay, type of travel etc. and needs to be discussed with a medical practitioner. For those countries with disease present, risk is highest in urban and semi-urban areas, but may also occur in rural areas; [insect avoidance measures](#) are highly recommended all year round. Travelvax believes that the best defence is to understand their habits, dress properly and use an [effective insect repellent](#) in the correct manner. Consult a medical practitioner for your specific risk.

### SCHISTOSOMIASIS

Swimming or bathing in freshwater is the main cause of infection. Avoid freshwater contact if you are unsure of origin.

## STIs

Low incidence of sexually transmitted and blood-borne diseases. Safe sex and needle precautions should be followed by all travellers.

## ALTITUDE

Altitude illness is a risk to travellers ascending higher than 2000 meters. Preventative medication is available. Consult a medical practitioner if ascending to 2000 metres or above.

## ADDITIONAL INFORMATION

### PERSONAL SAFETY

In order to check before and during travel for any high-risk areas visit the Smartraveller website. Avoid unnecessary displays of wealth or valuables and minimise the amounts of cash carried. Keep secure records of passport/credit card/licence numbers. For more safety tips visit: [www.smartraveller.gov.au](http://www.smartraveller.gov.au).

### MEDICAL CARE

Comprehensive medical facilities available only in the major population centres and/or specific private clinics. Sterile surgical products (syringes/needles/sutures) are not always available. Check for any contacts supplied by your emergency assistance organisation (nominated by your travel insurer) or with IAMAT (International Association of Medical Assistance for Travellers).

### FIRST AID KITS & ACCESSORIES

Carry a basic [first aid kit](#), customised to your activities. For longer stays consider a needle and syringe pack and medication for travellers diarrhoea.

**More information on Botswana is available during your pre-travel consultation with Travelvax.  
Call 1300 360 164 for the location of the clinic nearest to you.**