

RUSSIA

ABOUT RUSSIA VACCINATIONS

Russia stretches east from the Baltic Sea in Europe across North Asia to the Pacific Ocean. The climate varies from region to region in Russia. Northern Russia has harsh weather: January temperatures average -34°C. South-eastern Russia experiences a much milder climate, with an average temperature in July of 7°C. The temperature may reach 20°C in the western region near Moscow.

OVERALL RISK

Medium Risk - Preventable disease occur sporadically or at moderate levels. Consult a Travelvax clinic for your specific risk if travelling outside resorts, 5 star accommodation or for periods longer than a few days.

MALARIA RISK

No risk to travellers.

VACCINE PREVENTABLE DISEASES

YELLOW FEVER

No certificate required and no risk to travellers.

COVID-19

For the most up to date requirements for entry into Australia, go to Smartraveller.

Please click this COVID-19 Travel Regulations Map (IATA) to view the latest Russia COVID travel requirements.

HEPATITIS A

Moderate risk to travellers, vaccination recommended.

HEPATITIS B

Moderate presence of Hepatitis B in local population. Vaccination would be recommended.

TYPHOID

Moderate risk for most travellers. Vaccination recommended for travel to smaller cities, villages and rural areas outside usual tourist routes. Some medical conditions pre-dispose to infection; whether vaccinations would be recommended should be discussed with a medical practitioner. Consult a medical practitioner for your specific risk.

INFLUENZA

Northern hemisphere winter risk (November to April). Most common vaccine preventable illness in travellers. Vaccine recommended, effective for 1 year.

RABIES

Disease present. Recommendation for vaccination will depend on specific itinerary and activities planned. Generally rabies vaccination is advised for high risk individuals such as veterinarians or animal handlers, cavers. Additionally for higher risk travellers who plan: extended periods outdoors, rural travel, adventurous activities including bicycling; also expats or long-term travellers to endemic regions and children (risk of more severe or risk-prone bites and may not report contact at all).

JAPANESE ENCEPHALITIS

Disease present. Seasonal risk will vary by country. Whether vaccinations will be recommended will depend on itinerary, length of stay, type of travel etc and needs to be discussed with a medical practitioner. Risk is highest around pig farms and in agricultural areas. Mosquito avoidance measures are highly recommended all year round. Consult a medical practitioner for your specific risk, particularly if travelling during the wet season.

POLIO

No risk to travellers.

TICKBORNE ENCEPHALITIS

Vaccine is available in Australia - medical practitioners must apply through the special access scheme of the TGA. Recommended for high risk travellers: long-term residents, stays in rural and forested areas, campers and hikers, consuming unpasteurised dairy products. The disease is most active from April through August.

MENINGOCOCCAL MENINGITIS

Low risk to travellers.

CHOLERA

No risk to travellers.

OTHER DISEASES

TRAVELLERS DIARRHOEA

Moderate risk. Travellers' Diarrhoea affects 20-60% of overseas travellers, food and water precautions are essential. Self-treatment medications may be recommended.

INSECT-BORNE DISEASES

No risk to travellers.

SCHISTOSOMIASIS

No risk to travellers.

STIs

Low incidence of sexually transmitted and blood-borne diseases. Safe sex and needle precautions should be followed by all travellers.

AITITUDE

Altitude illness is a risk to travellers ascending higher than 2000 meters. Preventative medication is available. Consult a medical practitioner if ascending to 2000 metres or above.

ADDITIONAL INFORMATION

PERSONAL SAFETY

In order to check before and during travel for any high-risk areas visit the Smartraveller website. Avoid unnecessary displays of wealth or valuables and minimise the amounts of cash carried. Keep secure records of passport/credit card/licence numbers. For more safety tips visit: www.smartraveller.gov.au.

MEDICAL CARE

Limited medical facilities available. Unless travelling with a well-equipped organisation, a high level of self-sufficiency in terms of first aid kits and sterile equipment is recommended. An evacuation contingency should be a part of your travel insurance. Check for any contacts supplied by your emergency assistance organisation (nominated by your travel insurer) or with IAMAT (International Association of Medical Assistance for Travellers).

FIRST AID KITS & ACCESSORIES

Carry a basic <u>first aid kit</u>, customised to your activities. For longer stays consider a needle and syringe pack and medication for travellers diarrhoea.

More information on Russia is available during your pre-travel consultation with Travelvax. Call 1300 360 164 for the location of the clinic nearest to you.